

Low-density, one-dimensional quantum gases in a split trap

Th. Busch* and G. Huyet

Department of Physics, National University of Ireland, UCC, Cork, Ireland

(Dated: February 2, 2008)

We investigate degenerate quantum gases in one dimension trapped in a harmonic potential that is split in the centre by a pointlike potential. Since the single particle eigenfunctions of such a system are known for all strengths of the central potential, the dynamics for non-interacting fermionic gases and low-density, strongly interacting bosonic gases can be investigated exactly using the Fermi-Bose mapping theorem. We calculate the exact many-particle ground-state wave-functions for both particle species, investigate soliton-like solutions, and compare the bosonic system to the well-known physics of Bose gases described by the Gross-Pitaevskii equation. We also address the experimentally important questions of creation and detection of such states.

PACS numbers: 03.75.Fi, 05.30.Jp

I. INTRODUCTION

Low dimensional quantum gases are fastly progressing from being a solely successful theoretical concept [1, 2, 3, 4, 5] towards becoming experimentally available systems in the laboratories [6, 7, 8, 9, 10]. Small scale trapping potentials for low temperature samples allow the creation of degenerate fermionic or bosonic gases, for which the dynamics along two spatial directions can be 'frozen out'. Recent experimental progress has been the observation of quasi-one-dimensional Bose-condensates showing spatial phase fluctuations [11] and the generation of trapping frequencies in two-dimensional optical lattices of up to about 120kHz [12].

At very low temperatures the interaction between two neutral particles can often be described using a hard core potential that is dependent on a single parameter only, the s -wave scattering length. The Tonks-Girardeau (TG) gas is a model for a one-dimensional bosonic gas made from particles interacting via such a hard-core potential [3, 5] and in the limit of point-like interactions it can be solved exactly using the Fermi-Bose mapping theorem [3]. This theorem is one of the particularities of one-dimensional systems and it states that the properties of a strongly interacting, hard-core bosonic gas can be calculated by considering a non-interacting fermionic system. The first examples for which the TG gas model was solved were free space and box-like systems with periodic boundary conditions, and recently Wright and Girardeau have calculated its solution in an external harmonic potential [13, 14].

In order to find the many-particle solutions to a given geometry for the TG gas (or for non-interacting fermions) one must know the exact single particle eigenstates. However, since the list of exactly solvable single particle problems in quantum mechanics is limited, there are even fewer many-particle problems that can be exactly solved. Here we present a further solvable system, which com-

prises of a gas trapped in a harmonic oscillator potential split in the center by a point-like potential. This model can, on the one hand, be seen as a limit of double well situations [15], which have recently been very thoroughly investigated for examining weakly coupled Bose-Einstein condensates [16]. On the other hand it is a good approximation to the situation where an impurity is trapped in the center of a gas.

In Fermi-Bose mapping theory the bosonic wave-function can be calculated directly from an appropriately chosen fermionic one by symmetrization. Since the symmetry or antisymmetry of a wave function does not have an influence on the density distribution, the spatial density profiles for fermionic and bosonic samples in this limit are indistinguishable. Therefore, whenever results concerning density distributions are presented in this paper, they apply to fermionic as well as to bosonic samples.

The organization of the paper is the following: we will first briefly introduce the Fermi-Bose mapping theorem and its application to the TG gas. Then we will introduce the δ -split harmonic oscillator and present its exact solutions for arbitrary strength of the splitting potential. After this we will derive the exact many-particle ground-state wave-functions for the fermionic and the bosonic case in the limit of a completely split oscillator and investigate the dynamical behaviour when the central potential strength is changed. To compare the results to Bose gases in the Gross-Pitaevskii limit, we investigate a soliton-like solution. We suggest different methods for experimentally realizing a system like this and outline a strategy to observe it.

II. FERMİ-BOSE MAPPING THEOREM

We consider a gas of N neutral bosonic atoms of mass m at zero temperature in one dimension. The transverse dynamics is assumed to be 'frozen out' by a very stiff harmonic potential. The density of the gas shall be low enough to fulfill the requirement that the system is in the Tonks-Girardeau limit [17, 18]. In this limit the interaction between the particles can be treated as point-like.

*Electronic address: thbusch@phys.ucc.ie

Allowing that the atoms are longitudinally trapped in an external potential, $V(x)$, the many-particle Hamiltonian can be written as

$$\hat{H} = \sum_{i=1}^N \left[-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x_i^2} + V(x_i) \right] + g_1 \sum_{i<j}^N \delta(|x_i - x_j|). \quad (1)$$

Here the first sum represents the single particle Hamiltonian, $\hat{H}_0$, and the second sum accounts for the inter-particle interaction. The coupling constant is given by $g_1 = -2\hbar^2/2ma_{1D}$, with the repulsive, one-dimensional s -wave scattering length given by $a_{1D} = (-a_{\perp}^2/2a_{3D})[1 - Ca_{3D}/a_{\perp}]$ [17]. Here a_{3D} is the three-dimensional s -wave scattering length, $a_{\perp}$ the size of the ground state of the transversal harmonic trapping potential and $C = 1.4603\dots$. In the Tonks-Girardeau limit the one-dimensional coupling between the particles becomes large and we will consider $g_1 \rightarrow \infty$ in the Hamiltonian (1). In this situation, instead of including the scattering term into the Hamiltonian, it is more convenient to account for the strong repulsive interactions using a constraint on the allowed solutions [14]

$$\psi = 0 \quad \text{if} \quad |x_i - x_j| = 0, \quad i \neq j. \quad (2)$$

Since this constraint is equivalent to the demand of the Pauli exclusion principle for a gas of spin-less fermions, one can calculate the bosonic eigenfunctions, ψ_B , of the Hamiltonian of eq. (1) from the fermionic eigenfunctions, ψ_F , of the non-interacting Hamiltonian by proper symmetrisation

$$\Psi_B(x_1, \dots, x_N) = |\Psi_F(x_1, \dots, x_N)|. \quad (3)$$

This procedure is known as the Fermi-Bose mapping theorem [3, 4]. It can be generalized to time dependent problems.

To find the many-particle wave-function of the interacting Bose-gas we therefore need to solve the Slater determinant for non-interacting fermions. This determinant is composed of the eigenstates of the corresponding single particle Hamiltonian, $\hat{H}_0$. However, the number of physical systems where the single particle eigenstates are available analytically is very small. One example is the δ -split harmonic oscillator and we will review its single particle states in the next section.

III. EIGENSTATES OF THE δ -SPLIT HARMONIC TRAP

The single particle Hamiltonian describing a one-dimensional harmonic trap of frequency ω with a δ -potential of strength κ in the center is given by

$$\hat{H}_0 = -\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + \frac{1}{2}m\omega^2 x^2 + \kappa\delta(x). \quad (4)$$

A sketch of the potential is shown in the upper left of Fig. 1. One can note immediately that the odd eigenfunctions, ψ_{2n+1} , of the undisturbed harmonic oscillator ($\kappa = 0$) are still eigenfunctions of the Hamiltonian $\hat{H}_0$, since they are not affected by the potential ($\psi_{2n+1}(0) = 0$). In particular, they are independent of the strength κ of the δ -function.

To find the even eigenfunctions of the Hamiltonian in eq. (4), let us scale all quantities in units of the ground state size $a_0 = \sqrt{\hbar/2m\omega}$ and the energy $\epsilon_0 = \hbar\omega$ of the harmonic oscillator with $\kappa = 0$. The stationary Schrödinger equation then takes the form

$$\left(-\frac{d^2}{dx^2} + \frac{1}{4}x^2 + \tilde{\kappa}\delta(x) + \epsilon_n \right) \phi_n(x) = 0, \quad (5)$$

where we have also introduced the normalized energy $\epsilon_n = -E_n/(\hbar\omega)$ and the normalized delta function strength $\tilde{\kappa} = \kappa a_0/\epsilon_0$. Note the negative sign of the normalized energy, which has been chosen for easier mathematical treatment. One can see that for $x > 0$ eq. (5) is equal to the differential equation for the parabolic cylinder functions and the solutions that have the physically correct asymptotic behavior are Whittaker's functions, $U(\epsilon_n, x)$, given by [20]

$$U(\epsilon_n, x) = \cos\left(\frac{\pi}{4} + \frac{\pi\epsilon_n}{2}\right) Y_1 - \sin\left(\frac{\pi}{4} + \frac{\pi\epsilon_n}{2}\right) Y_2, \quad (6)$$

$$Y_1 = \frac{\Gamma\left(\frac{1}{4} - \frac{1}{2}\epsilon_n\right)}{\sqrt{\pi}2^{\left(\frac{1}{4} + \frac{1}{2}\epsilon_n\right)}} e^{-\frac{1}{4}x^2} M\left(\frac{1}{4} + \frac{1}{2}\epsilon_n, \frac{1}{2}, \frac{1}{2}x^2\right), \quad (7)$$

$$Y_2 = \frac{\Gamma\left(\frac{3}{4} - \frac{1}{2}\epsilon_n\right)}{\sqrt{\pi}2^{\left(-\frac{1}{4} + \frac{1}{2}\epsilon_n\right)}} e^{-\frac{1}{4}x^2} x M\left(\frac{3}{4} + \frac{1}{2}\epsilon_n, \frac{3}{2}, \frac{1}{2}x^2\right). \quad (8)$$

The $M(a, b, c)$ are the confluent hypergeometric functions. To first construct the symmetric eigenfunctions of the Hamiltonian in eq. (4) on the domain $x \neq 0$ we can now define [19]

$$\phi_n(x) = CU(\epsilon_n, |x|), \quad (9)$$

with the normalization constant C chosen such that the wave-function is normalized to one. Finally, evaluating the continuity condition at $\phi_n(x = 0)$

$$\frac{d}{dx}\phi_n(0^+) - \frac{d}{dx}\phi_n(0^-) = \tilde{\kappa}\phi_n(0), \quad (10)$$

we get an implicit relation between the strength of the δ -function and the eigenvalues of the Hamiltonian

$$\frac{\Gamma\left(\frac{3}{4} + \frac{1}{2}\epsilon_n\right)}{\Gamma\left(\frac{1}{4} + \frac{1}{2}\epsilon_n\right)} = -\tilde{\kappa}. \quad (11)$$

Since the Γ -function has poles for negative integer values, one can immediately see that the energy eigenvalues for a strong central barrier ($\kappa \rightarrow \infty$) converge to $E_n \rightarrow \frac{3}{2}\hbar\omega, \frac{7}{2}\hbar\omega, \frac{11}{2}\hbar\omega, \dots$. They become identical to the values of the odd eigenfunctions in the limit $\kappa = \infty$, leading to a double degeneracy of all eigenstates [19]. Fig. 1




FIG. 1: Sketch of the delta split harmonic oscillator and the respective single particle ground state ($n = 0$) densities for different values of κ . The functions for $\kappa = 0, 1, \infty$ correspond to the energy eigenvalues of $E_n = \frac{1}{2}\hbar\omega, 1.08\hbar\omega, \frac{3}{2}\hbar\omega$ respectively.

displays the square of the ground state eigenfunctions of the Hamiltonian in eq. (4) for different values of κ . One can clearly see the development of a node at $x = 0$ for increasing strength of the splitting potential.

In the limit $\kappa = \infty$, when the neighbouring even and odd eigenfunctions become degenerate, the Whittaker functions of the even states simplify to Hermite polynomials with an absolute value argument.

IV. QUANTUM GAS IN A δ -SPLIT HARMONICS TRAP

To find the exact many-particle wave-function for bosons or fermions, one has to calculate the Slater determinant from the single particle eigenfunctions $\phi_n(x)$ and, for the bosons, apply the symmetrization procedure to it. Due to the complicated structure of the Whittaker functions for finite values of κ this can only be done numerically. However, for the limiting case of an infinitely high central δ -function this can in fact be done analytically. The eigenfunctions are in this case given by

$$\psi_n(x) = C_n e^{-\frac{x^2}{2}} H_n(x) \quad \text{for } n \text{ odd}, \quad (12a)$$

$$\psi_n(x) = C_{n+1} e^{-\frac{|x|^2}{2}} H_{n+1}(|x|) \quad \text{for } n \text{ even}, \quad (12b)$$

with the normalization $C_n = (\sqrt{\pi}a_0 2^n n!)^{-\frac{1}{2}}$, and the Slater determinant is defined by

$$\psi_F(x_1, \dots, x_N) = \frac{1}{\sqrt{N!}} \det_{(n,j)=(0,1)}^{(N-1,N)} \psi_n(x_j). \quad (13)$$

Inserting the explicit expressions for the Hermite polynomials into the determinant and applying basic multi-

linear operations, the determinant can be reduced to

$$\det_{(n,j)=(0,1)}^{(N-1,N)} \psi_n(x_j) = C 2^{\frac{N^2}{2}} \left[\prod_j e^{-\frac{x_j^2}{2}} \right] \times \begin{vmatrix} |x_1| & x_1 & |x_1|^3 & x_1^3 & \dots & |x_1|^{n-1} & x_1^{n-1} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ |x_n| & x_n & |x_n|^3 & x_n^3 & \dots & |x_n|^{n-1} & x_n^{n-1} \end{vmatrix} \quad (14)$$

where $C = \prod_{n=1}^{N/2} C_{2n-1}^2$. Unfortunately in this form the determinant cannot be easily solved, due to the appearance of the absolute values.

We can, however, rewrite the problem in a more amenable form. It is easy to see that for the case of $\kappa = \infty$ due to the double degeneracy of each energy level, one can choose a new set of eigenfunctions given by $\phi_n^\pm = \frac{1}{\sqrt{2}}(\psi_{2n} \pm \psi_{2n+1})$ [15]. These eigenfunction correspond to the particles being physically located entirely either left or right of the barrier and the wave-functions $\phi_n^\pm$ are mutually symmetric with respect to $x = 0$. Therefore we can restrict our treatment to the spatial area $x > 0$ and consider only $\frac{N}{2}$ particles (for simplicity we assume an even number of particles). The Slater determinant can then be explicitly calculated by just employing the odd Hermite polynomials of eq. (12a)

$$\det_{(n,j)=(0,1)}^{(\frac{N}{2}-1, \frac{N}{2})} H_{2n+1}(x_j) = 2^{\frac{N^2}{8}} \left[\prod_j x_j \right] \prod_{(j,k)=(1,j+1)}^{(\frac{N}{2}, \frac{N}{2})} (x_j^2 - x_k^2) \quad (15)$$

which immediately gives a simple and exact expression for the fermionic many-particle wave-functions of eq. (13). The bosonic form then follows directly from eq. (3) and is given by

$$\Psi_B(x_1, \dots, x_N) = \frac{\tilde{C}}{\sqrt{N!}} 2^{\frac{N^2}{8}} \left[\prod_j e^{-\frac{x_j^2}{2}} |x_j| \right] \times \prod_{(j,k)=(1,j+1)}^{(N/2, N/2)} |x_j^2 - x_k^2| \quad (16)$$

with $\tilde{C} = \prod_{n=1}^{N/2} C_{2n-1}$. It resembles the wave-function found by Girardeau *et al.* [14] for the undisturbed harmonic oscillator, and differs due to the reason that we have constructed the Slater determinant only from the odd Hermite polynomials.

In Fig. 2 the density of a cloud of $N = 50$ bosons is shown for different heights of the central potential. One can clearly see that the central notch gets deeper with increasing values of κ , but one also finds an increase in density of about 25% just around the hole for large values of κ . As one would expect, the influence of the δ -potential only extends a short distance (the 'healing length') from $x = 0$ and does not have any influence on, say, the size of the atomic cloud.




FIG. 2: Ground state density for a one dimensional quantum gas of 50 particles in a δ -split harmonic trap. The strength of the δ -function is given by κ . Note the increase in density in the center of the trap around the notch. Note also that the appearance of the δ -function has no influence on the size of the atomic cloud.

The density at $x = 0$ of the gas as a function of κ is displayed in the LHS of Fig. 3. One can see that for values of κ of a few ω the density drops quickly towards zero. On the RHS the full-width-half-max size of the central notch is shown. As compared to the solitonic solutions for Bose gases described by the Gross-Pitaevskii equation where the width of the notch decreases with increasing darkness, the opposite behaviour is found here. This is of course due to the fact that the density void of the soliton in a Bose condensate is due to a phase jump in the wave-function and not an external potential.

V. DARK SOLITONS

As already mentioned in the last section, the density distributions shown in Fig. 2 for finite κ resemble the dark and grey soliton states of the Gross-Pitaevskii equation (GPE). Dark (and grey) solitons are solutions to the GPE




FIG. 3: The left hand side shows the density of a gas at $x = 0$ depending on the strength of the delta function. The right hand side shows the FWHM of the central notch. Note that the width increase with increasing 'darkness'.

with repulsive interaction and their existence in dilute Bose-condensates has been experimentally demonstrated [21, 22]. Recently also the appearance of dark solitons in TG gas has been examined in a toroidal trap [23].

Having derived the exact many-particle state for the pierced harmonic oscillator, we can now compare the TG 'notch' state with soliton solutions of the Gross-Pitaevskii equation in a trap. Note, that we have not only the 'dark notch' solution at hand, but we also know the full family of solutions for 'grey notches', where the density is reduced but does not go to zero.

We are, however, again not able to derive any analytical expression for the grey soliton densities, since the summations over the eigenstate densities involve a summation over ϵ_n . This doesn't represent a series of integers nor is there an easy relationship between the integer n and the ϵ_n and we therefore resort to numerical simulations.

To simulate the time evolution of the bosonic gas, one can make use of a time dependent version of the Fermi-Bose mapping theorem [23]. The many-particle fermionic wave-function is then given by

$$\psi_F(x_1, \dots, x_N; t) = C \det_{(i,j)=(0,1)}^{(N-1,N)} \phi_i(x_j, t), \quad (17)$$

and the single particle eigenstates of the external potential evolve under the time-dependent Schrödinger equation

$$i\hbar \frac{\partial \phi_i(x, t)}{\partial t} = \left[-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} + V(x, t) \right] \phi_i(x, t). \quad (18)$$

Symmetrization of these states then leads to the appropriate solutions for the bosonic case. To investigate the dynamics of the notch, we assume that we can prepare the system in the stationary state we want for fixed κ and then we instantaneously switch $\kappa = 0$.

First, it is easy to understand that the recurrence frequency for the density notch in the gas is given by the trap frequency. Since the density distribution is in particular the density distribution for non-interacting fermions, after one trapping period ω every fermion is back in the same state where it started and the notch is recreated (see Fig. 4). Even though this result is to be expected for non-interacting fermions, it is not at all obvious for a system of strongly interacting bosons. In particular it is in contrast to behaviour of dark solitons in the GPE, for which the oscillation frequency is given by $\omega/\sqrt{2}$ [24].

Since the filling in and the oscillation of the dark notch as seen in Fig. 4 are purely due to the even eigenfunctions (the odd ones are stationary states for any value of κ), one could consider to stabilize the density notch by imprinting a phase-step with $\Delta\Phi = \pi$ onto the sample. This would artificially create the same stabilization mechanism that the soliton solution of the GPE has, i.e. it would transform the single particle states with an even parity to states with an odd one. However, since one cannot make this process parity selective, at the same




FIG. 4: Density at $x = 0$ during one oscillation period after κ has instantaneously been changed from $\kappa = \infty$ to $\kappa = 0$. One can clearly see the recurrence of the notch after one trap period.

time it would transfer the odd states into even states and nothing would be won.

VI. DETECTION

The absence of a self-similar free expansion has been pointed out as one criterion to distinguish the TG gas from a one-dimensional Bose-gas in the Gross-Pitaevskii regime [25]. Since the atomic cloud is only disturbed on a very small length scale by the central δ -function, and, as can be seen from Fig. 4, the lifetime for the density notch once the central barrier is removed is much shorter than a trap frequency, the modification due to the split trap has no measurable effect on the size and dynamics of the expanding cloud (see Fig. 5). Especially, since the lifetime of the density notch is so short, one cannot observe any broadening of it

This fact can also be seen by realizing that the energy increase for finite κ is at maximum $\Delta E \rightarrow \frac{N}{2}\hbar\omega$ for the limiting case of $\kappa \rightarrow \infty$, which is very small on the scale of the total energy of the cloud, $E_{tot} = N^2/2$. (Note that the chemical potential alone is already $\mu = (N-1/2)\hbar\omega$.)

We suggest here a different approach that relies on the insensibility of the state towards phase imprinting. In standard traps ($\kappa = 0$) phase imprinting has a substantial effect on the density distribution of a sample, a fact which has been successfully utilized for creating dark solitons in ground state Bose-Einstein condensates. In the dynamics following the imprinting the gas tries to adjust its density distribution to the phase, thereby excit-




FIG. 5: Integrated absolute difference between the density distributions of the ground state gas for $\kappa = 0$ and $\kappa = \infty$ during the free expansion. The fast decay is a sign of the indistinguishability of the two ground states via the density distributions. The inset shows the density $\rho(x=0)$ for $\kappa = \infty$ (broken line) and for $\kappa = 0$ (full line). One can see that the hole vanishes at very early stages during the free evolution. Due to the absence of an external potential, there are no recurrences here. The timescale in both graphs is in units of the inverse trap frequency.

ing various phonon modes and modifying the Thomas-Fermi density distribution. Recently phase imprinting was theoretically investigated for fermionic samples, and the emergence of solitonic structures was also found there [26]. However, since the phase evolution of the single particles in a Fermi sea is not coherent, the effects found were less pronounced.

To demonstrate the insensibility of the bosonic and fermionic states in a δ -split trap, we consider here the limiting case of $\kappa \rightarrow \infty$, and the reaction to an imprinted phase of $\Delta\phi = \pi$. Applying this imprinting only spatially selectively for, say, $x < 0$, one can see immediately that the eigenfunctions $\psi_n(x)$ with odd values of n turn into $\psi_{n-1}(|x|)$ and vice versa. Since after the imprinting we again deal only with stationary eigenfunctions, the phase imprinting (assuming an ideal imprinting mask) does not have any observable effect on the density distribution. Mathematically one can see immediately that the Slater determinant due to its multilinearity is unchanged by the imprinting of the phase (for an even number of particles). Therefore the absence of distortions could be used as a possible tool to detect the split trap state.

VII. EXPERIMENTAL REALIZATION OF THE δ -POTENTIAL

Let us finally remark on the experimental realization of such a potential, since the δ -function is usually regarded

as a mathematical idealization far from experimental reality. A very successful method to manipulate the density profile of an atomic cloud is to employ dipole forces with appropriately detuned lasers. It is nevertheless experimentally quite challenging to focus and stabilize a laser beam to such a small diameter that it represents a point-like potential to the atoms as necessary for the situation considered here. However, since low densities are best reached in shallow traps, the ratio of the size of the single particle states vs. the fwhm of the focused laser beam increases with increasing trap size, thereby creating a more and more localized potential. The tuning of κ could in this case be done 'easily' by adjusting the intensity of the laser beam.

Here we suggest an alternative method which at the same time also allows for easy tuning of κ , even though it is also experimentally more challenging.

If optical potentials do not suffice in terms of localization, the obvious next step is to employ potentials made from matter waves. Multicomponent condensates have been investigated in the last years and the conditions for such systems to exhibit phase separation have been determined. Depending on the internal energy of the atoms and on their respective masses shell like structures have been predicted, where one component occupies the inner part of the trap and the other component occupies the outer part. By realizing a situation like this and then reducing the number of atoms occupying the central part of the trap, one can, in the ideal case, come to the situation where a single atom is trapped in the trap center and the second component is arranged around it. This would correspond to an almost perfect point-like potential for the atoms of the outer component. With the help of a selective Feshbach resonance the strength of the interaction between the single atom and the gas could then be adjusted.

VIII. CONCLUSIONS

We have investigated the behaviour of low density, one-dimensional Bose and Fermi gases in a δ -split har-

monic trap. We have derived the exact many-particle wave function for non-interacting fermions and were able, due to the Fermi-Bose mapping theorem, to determine also the exact expression for the many-particle wave function for the bosonic TG gas in the limiting case of an infinitely high δ -barrier. Both many-particle wave functions turned out to be simple products over quadratic terms. We have thereby presented another of the rare examples of exactly solvable models for many-particle systems.

We have further numerically investigated the dynamical behavior of many-particle, split-trap eigenstate when the central barrier is suddenly removed and compared it to the dynamics of the dark soliton solutions in the GPE. The recurrence time for the 'notch' in the linear low density limit was found to be the trapping frequency, which is different to the oscillation frequency of the dark Gross-Pitaevskii soliton of $\omega/\sqrt{2}$.

Finally, we have suggested using the systems insensitivity to phase imprinting as a tool for detecting the presence of the notch-state and employing a second component and a selective Feshbach resonance to create and manipulate the point-like central potential.

Acknowledgments

This work was financially supported by the European Union under contract3 HPRN-CT-2000-00034-VISTA, by the Science Foundation Ireland under the grant sfi/01/fi/co and by the Royal Irish Academy. Discussions with S. Nic Chormaic and J. Denschlag are gratefully acknowledged. TB would like to thank John McNerney for his support.

-
- [1] E.H. Lieb and W. Liniger, Phys. Rev. **130**, 1605 (1963).
 - [2] E.H. Lieb, Phys. Rev. **130**, 1616(1963)
 - [3] M. Girardeau, J. Math. Phys. **1**, 516 (1960).
 - [4] M. Girardeau, J. Math. Phys. (N.Y.) **1**, 516 (1960); Phys. Rev. **139**, B500 (1965).
 - [5] L. Tonks, Phys. Rev. **50**, 955 (1936).
 - [6] A. Görlitz, J.M. Vogels, A.E. Leanhardt, C. Raman, T.L. Gustavson, J.R. Abo-Shaer, A.P. Chikkatur, S. Gupta, S. Inouye, T. Rosenband and W. Ketterle, Phys. Rev. Lett. **87**, 130402 (2001).
 - [7] R. Dumke, T. Muther, M. Volk, W. Ertmer, and G. Birkel, Phys. Rev. Lett. **89**, 220402 (2002).
 - [8] N.H. Dekker, C.S. Lee, V. Lorent, J.H. Thywissen, S.P. Smith, M. Drndic, R.M. Westervelt, and M. Prentiss, Phys. Rev. Lett. **84** 1124 (1999).
 - [9] H. Ott, J. Fortagh, G. Schlotterbeck, A. Grossmann, and C. Zimmermann, Phys. Rev. Lett. **87**, 230401 (2001).
 - [10] B.K. Teo and G. Raithel, Phys. Rev. A **65**, 051401(R) (2002).
 - [11] S. Dettmer, D. Hellweg, P. Rytty, J.J. Arlt, W. Ertmer, K. Sengstock, D.S. Petrov, G.V. Shlyapnikov, H. Kreutzmann, L. Santos, and M. Lewenstein, Phys. Rev. Lett. **87**, 160406 (2001).
 - [12] M. Greiner, I. Bloch, O. Mandel, T.W. Hnsch, and T. Esslinger, Phys. Rev. Lett. **87**, 160405 (2001).
 - [13] M.D. Girardeau and E.M. Wright, Phys. Rev. Lett. **84**,

- 5239 (2000)
- [14] M.D. Girardeau, E.M. Wright, and J.M. Triscari, Phys. Rev. A **63**, 033601 (2001).
 - [15] J. Javanainen and M. Wilkens, Phys. Rev. Lett. **25**, 4675 (1997).
 - [16] A. Smerzi, S. Fantoni, S. Giovanazzi, S.R. Shenoy, Phys. Rev. Lett. **79**, 4950 (1997).
 - [17] M. Olshanii, Phys. Rev. Lett. **81**, 938 (1998).
 - [18] D.S. Petrov, G.V. Shlyapnikov, and J.T.M. Walraven, Phys. Rev. Lett. **85**, 3745 (2000).
 - [19] Th. Busch, B.G. Englert, K. Rzazewski, and M. Wilkens, Found. of Phys. **28**, 549 (1998).
 - [20] M. Abramowitz and I.A. Stegun, Eds., *Handbook of Mathematical Functions* (Dover, New York, 1972).
 - [21] S. Burger, K. Bongs, S. Dettmer, W. Ertmer, K. Sengstock, A. Sanpera, G.V. Shlyapnikov, and M. Lewenstein, Phys. Rev. Lett. **83**, 5198 (1999).
 - [22] J. Denschlag, J.E. Simsarian, D.L. Feder, C.W. Clark, L.A. Collins, J. Cubizolles, L. Deng, E.W. Hagley, K. Helmerson, W.P. Reinhardt, S.L. Rolston, B.I. Schneider, and W.D. Phillips, Science **287**, 97 (2000).
 - [23] M.D. Girardeau and E.M. Wright, Phys. Rev. Lett. **84**, 5691 (2000).
 - [24] Th. Busch and J.R. Anglin, Phys. Rev. Lett. **84**, 2298 (2000).
 - [25] P. Öhberg and L. Santos, cond-mat/0204611 (2002).
 - [26] T. Karpiuk, M. Brewczyk, L. Dobrek, M. A. Baranov, M. Lewenstein, and K. Rzazewski, Phys. Rev. A **66**, 023612 (2002).